

Relationships between the Phase Retrieval Problem and Permutation Invariant Embeddings

Radu Balan and Efstratios Tsoukanis

Department of Mathematics, University of Maryland, College Park, MD 20742

Emails: rvbalan@umd.edu , etsoukan@umd.edu

Abstract—This paper discusses the connection between the phase retrieval problem and permutation invariant embeddings. We show that the real phase retrieval problem for $\mathbb{R}^d/O(1)$ is equivalent to Euclidean embeddings of the quotient space $\mathbb{R}^{2 \times d}/S_2$ performed by the sorting encoder introduced in an earlier work. In addition, this relationship provides us with inversion algorithms of the orbits induced by the group of permutation matrices.

I. INTRODUCTION

The phase retrieval problem has a long and illustrious history involving several Nobel prizes along the way. The issue of reconstruction from magnitude of frame coefficients is related to a significant number of problems that appear in separate areas of science and engineering. Here is an incomplete list of some of these applications and reference papers: crystallography [1], [2], [3]; ptychography [4], [5]; source separation and inverse problems [6], [7]; optical data processing [8]; mutually unbiased bases [9], [10], quantum state tomography [11], [12]; low-rank matrix completion problem [13], [14]; tensor algebra and systems of multivariate polynomial equations [15], [16], [17]; signal generating models [18], [19], bandlimited functions [20], [21], radar ambiguity problem [22], [23], learning and scattering networks [24], [25], [26].

In [27], this problem was shown to be a special form of the following setup. Let H denote a real or complex vector space and let $A = \{a_i\}_{i \in I}$ be a frame for H . The phase retrieval problem asks whether the map $H \ni x \mapsto \alpha_A(x) = \{|\langle x, a_i \rangle|\}_{i \in I} \in l^2(I)$ determines x uniquely up to a unimodular scalar.

In this paper we focus on the finite dimensional real case of this problem (see also [28]), namely when $H = \mathbb{R}^d$. In this case, a frame $\mathcal{A} = \{a_1, \dots, a_D\} \subset \mathbb{R}^d$ is simply a spanning set. The group $O(1) = \{-1, +1\}$ acts on H by scalar multiplication. Let $\hat{H} = H/O(1)$ denote the quotient space induced by this action, where the equivalence classes (orbits) are

$$[x] = \{x, -x\}, \text{ for } x \neq 0, \quad [x] = \{0\}, \text{ for } x = 0.$$

The analysis operator for this frame is

$$T_A : H \rightarrow \mathbb{R}^D, \quad T_A(x) = (\langle x, a_k \rangle)_{k=1}^D. \quad (1)$$

The relevant nonlinear map α_A is given by taking the absolute value of entries of T_A :

$$\alpha_A : H \rightarrow \mathbb{R}^D, \quad \alpha_A(x) = (|\langle x, a_k \rangle|)_{k=1}^D. \quad (2)$$

Notice α_A produces a well-defined map on \hat{H} , which, with a slight abuse, but for simplicity of notation, will be denoted also by α_A . Thus $\alpha_A([x]) = \alpha_A(x)$.

Another customary notation that is often employed: a frame is given either as an indexed set of vectors, $\mathcal{A} = \{a_1, \dots, a_D\}$, or through the columns of a $d \times D$ matrix A . The matrix notation is not canonical, but this is not an issue here. We always identify $H = \mathbb{R}^d$ with its column vector representation in its canonical basis.

Definition 1. We say that (the columns of a matrix) $A \in \mathbb{R}^{d \times D}$ form/is a phase retrievable frame, if $\alpha_A : \hat{\mathbb{R}}^d \rightarrow \mathbb{R}^D$, $\alpha_A(x) = (|\langle x, a_k \rangle|)_{k=1}^D$ is an injective map (on the quotient space).

In a different line of works [29], [30], [31], [32] it

was recognized that the phase retrieval problem is a special case of Euclidean representations of metric spaces of orbits defined by certain unitary group actions on Hilbert spaces. Specifically, the setup is as follows. Let V denote a Hilbert space, and let G be a group acting unitarily on V . Let $\hat{V} = V/G$ denote the metric space of orbits, where the quotient space is induced by the equivalence relation $x, y \in V$, $x \sim y$ iff $y = g.x$, for some $g \in G$. Here $g.x$ represents the action of the group element $g \in G$ on vector x . For the purposes of this paper we specialize to the finite dimensional real case, $V = \mathbb{R}^{n \times d}$ and $G = S_n$, is the group of $n \times n$ permutation matrices acting on V by *left multiplication*. Other cases are discussed in aforementioned papers. In particular, in [30] the authors have shown a deep connection to graph deep learning problems. In [31], the authors linked this framework to certain graph matching problems and more. The bi-Lipschitz Euclidean embedding problem for the finite dimensional case is as follows. Given $\hat{V} = V/G$, construct a map $\beta : V \rightarrow \mathbb{R}^m$ so that, (i) $\beta(g.x) = \beta(x)$ for all $g \in G$, $x \in V$, and (ii) for some $0 < A \leq B < \infty$, and for all $x, y \in V$,

$$A \mathbf{d}([x], [y]) \leq \|\beta(x) - \beta(y)\| \leq B \mathbf{d}([x], [y]) \quad (3)$$

where $\mathbf{d}([x], [y]) = \inf_{g \in G} \|x - g.y\|_V$ is the *natural metric* on the quotient space \hat{V} .

In [30] the following embedding was introduced. Let $A \in \mathbb{R}^{d \times D}$ be a fixed matrix (termed as *key*) whose columns are denoted by a_1, \dots, a_D . The induced encoder $\beta_A : V \rightarrow \mathbb{R}^{n \times D}$ is defined by

$$\beta_A(X) = \downarrow(XA) = \begin{bmatrix} \Pi_1 X a_1 & \cdots & \Pi_D X a_D \end{bmatrix} \quad (4)$$

where $\Pi_k \in S_n$ is the permutation matrix that sorts in decreasing order the vector $X a_k$. It was shown in [30] that, for D large enough, β_A provides a bi-Lipschitz Euclidean embedding of \hat{V} . This motivates the following definition.

Definition 2. We say that $A \in \mathbb{R}^{d \times D}$ is a universal key for $\mathbb{R}^{n \times d}$ if $\beta_A : \widehat{\mathbb{R}^{n \times d}} \rightarrow \mathbb{R}^{n \times D}$, $\beta_A(X) = \downarrow(XA)$ is an injective map (on the quotient space).

The purpose of this paper is to show the equivalence between the real phase retrieval problem,

specifically the embedding α_A , and the permutation invariant embedding β_A defined above, in the special case $n = 2$.

II. MAIN RESULTS

Recall the Hilbert spaces $H = \mathbb{R}^d$ and $V = \mathbb{R}^{2 \times d}$. For $A \in \mathbb{R}^{d \times D}$ recall also the encoders $\alpha_A : \hat{H} \rightarrow \mathbb{R}^D$ and $\beta_A : \hat{V} \rightarrow \mathbb{R}^{2 \times D}$ given respectively by $\alpha_A(x) = (\langle x, a_k \rangle)_{k \in [D]}$, and $\beta_A(X) = \downarrow(XA)$. Our main result reads as follows.

Theorem 3. *In the case $n = 2$, the following are equivalent.*

- 1) α_A is injective, hence the columns of A form a phase retrievable frame;
- 2) β_A is injective, hence A is a universal key.

Remark 4. *Perhaps it is not surprising that, if an equivalence between the phase retrieval problem and permutation invariant representations is possible, then this should occur for $n = 2$. This statement is suggested by the observation that $O(1)$ is isomorphic with S_2 , the group of the 2×2 permutation matrices. What is surprising that, in fact, the two embeddings are intimately related, as the proof and corollaries show.*

Proof of Theorem 3. Let $X \in V = \mathbb{R}^{2 \times d}$. Denote by $x_1, x_2 \in \mathbb{R}^d$ its two rows transposed, that is

$$X = \begin{bmatrix} x_1^T \\ x_2^T \end{bmatrix}.$$

Notice that, for each $k \in [D]$, the k^{th} column of $\beta_A(X)$ is given by

$$\downarrow(X a_k) = \begin{bmatrix} \max(\langle x_1, a_k \rangle, \langle x_2, a_k \rangle) \\ \min(\langle x_1, a_k \rangle, \langle x_2, a_k \rangle) \end{bmatrix}.$$

The *key observations* are the following relationships between *min*, *max*, and the absolute value $|\cdot|$:

$$\begin{aligned} |u - v| &= \max(u, v) - \min(u, v) \\ u + v &= \max(u, v) + \min(u, v) \\ \max(u, v) &= \frac{1}{2}(u + v + |u - v|) \\ \min(u, v) &= \frac{1}{2}(u + v - |u - v|) \\ ||u| - |v|| &= \min(|u - v|, |u + v|) \end{aligned}$$

In particular, these show that: □

$$\begin{aligned} \begin{bmatrix} 1 & -1 \\ 1 & 1 \end{bmatrix} \beta_A(X) &= \begin{bmatrix} 1 & -1 \\ 1 & 1 \end{bmatrix} \cdot \downarrow(XA) = \\ &= \begin{bmatrix} |\langle x_1 - x_2, a_1 \rangle|, & \dots, & |\langle x_1 - x_2, a_D \rangle| \\ \langle x_1 + x_2, a_1 \rangle, & \dots, & \langle x_1 - x_2, a_D \rangle \end{bmatrix} \\ &= \begin{bmatrix} (\alpha_A(x_1 - x_2))^T \\ (T_A(x_1 + x_2))^T \end{bmatrix} \end{aligned}$$

Where, T_A was introduced in equation (1).

(1) \rightarrow (2) : Suppose that α_A is injective. Let $X = \begin{bmatrix} x_1^T \\ x_2^T \end{bmatrix}$ and $Y = \begin{bmatrix} y_1^T \\ y_2^T \end{bmatrix}$, such that $\beta_A(X) = \beta_A(Y)$. Then

$$\begin{aligned} \begin{bmatrix} 1 & -1 \\ 1 & 1 \end{bmatrix} \beta_A(X) &= \begin{bmatrix} 1 & -1 \\ 1 & 1 \end{bmatrix} \beta_A(Y) \\ \implies \begin{bmatrix} (\alpha_A(x_1 - x_2))^T \\ T_A(x_1 + x_2)^T \end{bmatrix} &= \begin{bmatrix} (\alpha_A(y_1 - y_2))^T \\ T_A(y_1 + y_2)^T \end{bmatrix}. \end{aligned}$$

But now, $\alpha_A(x_1 - x_2) = \alpha_A(y_1 - y_2) \implies x_1 - x_2 = y_1 - y_2$ or $x_1 - x_2 = y_2 - y_1$ and

$$T_A(x_1 + x_2) = T_A(y_1 + y_2)^T \implies x_1 + x_2 = y_1 + y_2$$

Thus we have that

$$\left\{ \begin{array}{l} x_1 = y_1 \\ x_2 = y_2 \end{array} \right\} \text{ or } \left\{ \begin{array}{l} x_1 = y_2 \\ x_2 = y_1 \end{array} \right\}$$

Either case means

$$\begin{aligned} \iff X = Y \text{ or } X &= \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} Y \\ \iff [X] &= [Y] \end{aligned}$$

So, β_A is injective.

(2) \rightarrow (1) : Suppose that β_A is injective. Let $x, y \in \mathbb{R}^d$ such that $\alpha_A(x) = \alpha_A(y)$, i.e.

$$|\langle x, a_k \rangle| = |\langle y, a_k \rangle|, \forall k \in [D]. \text{ Let } X = \begin{bmatrix} x^T \\ -x^T \end{bmatrix}$$

and $Y = \begin{bmatrix} y^T \\ -y^T \end{bmatrix}$. Then,

$$\begin{bmatrix} 1 & -1 \\ 1 & 1 \end{bmatrix} \beta_A(X) = \begin{bmatrix} \alpha_A(2x)^T \\ T_A(0)^T \end{bmatrix} = 2 \begin{bmatrix} \alpha_A(2x)^T \\ 0 \end{bmatrix}$$

and

$$\begin{bmatrix} 1 & -1 \\ 1 & 1 \end{bmatrix} \beta_A(Y) = \begin{bmatrix} \alpha_A(2y)^T \\ T_A(0)^T \end{bmatrix} = 2 \begin{bmatrix} \alpha_A(2y)^T \\ 0 \end{bmatrix}$$

Thus $\beta_A(X) = \beta_A(Y)$. Since β_A is assumed injective, it follows that $X = Y$ or $X = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} Y$. So, $x = y$ or $x = -y$. We conclude that $[x] = [y]$, so α_A is injective.

Corollary 5. If β_A is injective, then $D \geq 2d - 1$.

Corollary 6. If $D = 2d - 1$, then β_A is injective if and only if A is a full spark frame.

Both results follow necessary and sufficient conditions established in, e.g. [27]. Recall that a frame in \mathbb{R}^d is said *full spark* if any subset of d vectors is linearly independent (hence basis).

Remark 7. Assume $D = 2d - 1$. Note the embedding dimension for $\hat{V} = \widehat{\mathbb{R}^{2 \times d}}$ is $m = 2(2d - 1) = 4d - 2 = 2 \dim(V) - 2$. In particular this shows the minimal dimension of bi-Lipschitz Euclidean embeddings may be smaller than twice the intrinsic dimension of the Hilbert space where the group acts on. Both papers [30] and [31] present (bi)Lipschitz embeddings into $\mathbb{R}^{2 \dim(V)}$.

Remark 8. As was derived in the proof, α_A , β_A and T_A are intimately related:

$$\beta_A \left(\begin{bmatrix} x_1^T \\ x_2^T \end{bmatrix} \right) = \frac{1}{2} \begin{bmatrix} 1 & 1 \\ -1 & 1 \end{bmatrix} \begin{bmatrix} \alpha_A(x_1 - x_2)^T \\ T_A(x_1 + x_2)^T \end{bmatrix} \quad (5)$$

In particular, any algorithm for solving the phase retrieval problem solves also the inversion problem for β_A . Let $\omega_A : \mathbb{R}^D \rightarrow \mathbb{R}^d$ denote a left inverse of α_A on the metric space \mathbb{R}^d . This means $\omega_A(\alpha_A(x)) \sim x$ in $\mathbb{R}^d/O(1)$. Denote by T_A^\dagger a left inverse of the analysis operator (e.g., the synthesis operator associated to the canonical dual frame). Thus $T_A^\dagger T_A = I_d$. Then an inverse for β_A is:

$$\beta_A^{-1}(Y) = \frac{1}{2} \begin{bmatrix} T_A^\dagger(y_2) + \omega_A(y_1) \\ T_A^\dagger(y_2) - \omega_A(y_1) \end{bmatrix} \quad (6)$$

where $Y = \begin{bmatrix} y_1^T \\ y_2^T \end{bmatrix}$.

Remark 9. Equations (6) suggest a lower dimensional embedding than β_A . Specifically, first we compute the average $y_1 = \frac{1}{2}(x_1 + x_2)$ which is of size \mathbb{R}^d , and then encode the difference $x_1 - x_2$ using α_A , $y_2 = \alpha_A(x_1 - x_2)$. We obtain the following modified encoder, $\tilde{\beta}_A : \mathbb{R}^{2 \times d} \rightarrow \mathbb{R}^{d+D}$:

$$\tilde{\beta}_A(x) = \begin{bmatrix} \frac{1}{2}(x_1 + x_2)^T & \alpha_A(x_1 - x_2)^T \end{bmatrix}. \quad (7)$$

With the ω_A left inverse of α_A , the inverse of $\tilde{\beta}_A$

is given by:

$$\tilde{\beta}_A^{-1}(Y) = \begin{bmatrix} y_1 + \frac{1}{2}\omega_A(y_2) \\ y_1 - \frac{1}{2}\omega_A(y_2) \end{bmatrix} \quad (8)$$

where $y_1 = Y(1:d)$ and $y_2 = Y(d+1:d+D)$. In the case when $D = D_{\min} = 2d - 1$, the minimal embedding dimension is $m = d + D = 3d - 1$ (instead of $4d - 2$ or $4d = 2 \dim(V)$).

Reference [30] shows that an upper Lipschitz bound for embedding β_A is $\sigma_1(A)$, where $\sigma_1(A)$ is the largest singular value of A . Same reference shows that if β_A is injective then there is also a strictly positive lower Lipschitz bound without providing a formula. Using Equation (5) we provide explicit estimates of these bounds.

Theorem 10. *Assume $A \in \mathbb{R}^{d \times D}$ is a universal key for $\mathbb{R}^{2 \times d}$ (i.e., $\beta_A : \widehat{\mathbb{R}^{2 \times d}} \rightarrow \mathbb{R}^{2 \times D}$ is injective), or, equivalently (according to Theorem 3), the columns of A form a phase retrievable frame in \mathbb{R}^d (i.e., $\alpha_A : \widehat{\mathbb{R}^d} \rightarrow \mathbb{R}^D$ is injective). Then both α_A and β_A are bi-Lipschitz with same Lipschitz constants, where distances are given by $\mathbf{d}_{\text{PR}}([x], [y]) = \min(\|x - y\|, \|x + y\|)$ on \hat{H} , and $\mathbf{d}([X], [Y]) = \min_{P \in S_2} \|X - PY\|$ on \hat{V} , respectively. The optimal lower and upper Lipschitz constants are given by:*

$$A_0 = \min_{I \subset [D]} \sqrt{\sigma_d^2(A[I]) + \sigma_d^2(A[I^c])}, \quad B_0 = \sigma_1(A) \quad (9)$$

where $\sigma_1(A)$ is the largest singular value of A (equals the square-root of upper frame bound) and $\sigma_d(A[J])$ is the d^{th} singular value of submatrix of A indexed by J . Furthermore, these bounds are achieved by the following vectors. Let I_0 denote a optimal partition in (9) and let u_1, u_2 denote the normalized left singular vectors of $A[I_0]$ and $A[I_0^c]$, respectively, each associated to the d^{th} singular value. Let u be the normalized principal left singular vector associated to A . (i.e., associated to the largest singular value). Then:

- 1) The upper Lipschitz constant B_0 is achieved as follows: (i) for map α_A by vectors $x_{\max} = u$ and $y_{\max} = 0$; (ii) for map β_A by vectors $X_{\max} = \begin{bmatrix} u^T \\ 0 \end{bmatrix}$ and $Y_{\max} = 0$.
- 2) The lower Lipschitz constant A_0 is achieved as follows: (i) for map α_A by vectors $x_{\min} = u_1 + u_2$ and $y_{\min} = u_1 - u_2$; (ii) for map

$$\beta_A \text{ by vectors } X_{\min} = \begin{bmatrix} (u_1 + u_2)^T \\ 0 \end{bmatrix} \text{ and } Y_{\min} = \begin{bmatrix} u_1^T \\ u_2^T \end{bmatrix}.$$

Remark 11. *The optimal Lipschitz constants for the map α_A were obtained in [33], [34], including the optimizers. However, for reader's convenience, we prefer to give direct proofs of these results.*

Proof. 1) Upper Lipschitz constants.

- (i) Let $x, y \in \mathbb{R}^d$. Then

$$\begin{aligned} \|\alpha_A(x) - \alpha_A(y)\|^2 &= \sum_{i=1}^D \left| \langle a_i, x \rangle - \langle a_i, y \rangle \right|^2 = \\ &= \sum_{i=1}^D \min(|\langle a_i, x-y \rangle|^2, |\langle a_i, x+y \rangle|^2) \leq \\ &\leq \min\left(\sum_{i=1}^D |\langle a_i, x-y \rangle|^2, \sum_{i=1}^D |\langle a_i, x+y \rangle|^2\right) \\ &\leq \sigma_1^2(A) \mathbf{d}_{\text{PR}}([x], [y])^2. \end{aligned}$$

So $\sigma_1(A)$ is an upper Lipschitz bound for the map α_A . Now for $x_{\max} = u, y_{\max} = 0$ notice that

$$\begin{aligned} \|\alpha(x_{\max}) - \alpha(y_{\max})\|^2 &= \sum_{i=1}^D |\langle a_i, u \rangle|^2 = \\ &= \sigma_1^2(A) \|u\|^2 = \sigma_1^2(A) \mathbf{d}_{\text{PR}}([x_{\max}], [y_{\max}])^2. \end{aligned}$$

Thus, the upper Lipschitz constant $\sigma_1(A)$ is in fact optimal (tight).

- (ii) Map β_A . Let $X, Y \in \mathbb{R}^{2 \times D}$ and $P_0 \in S_2$ be a permutation that achieves the distance between X and Y , i.e. $\|X - P_0 Y\| = \mathbf{d}([X], [Y])$. Note that

$$\begin{aligned} \|\beta_A(X) - \beta_A(Y)\|^2 &= \sum_{k=1}^D \|(\Pi_k X - \Xi_k Y)_{a_k}\|^2 = \\ &= \sum_{k=1}^D \|(\Xi_k^T \Pi_k X - Y)_{a_k}\|^2 \end{aligned}$$

for some $\Pi_k, \Xi_k \in S_2$ that align the vectors. From rearrangement lemma we have that

$$\|(\Pi_k X - \Xi_k Y)_{a_k}\| \leq \|(X - P_0 Y)_{a_k}\|, \quad \forall k \in [D]$$

so,

$$\begin{aligned} \sum_{k=1}^D \|(\Xi_k^T \Pi_k X - Y)_{a_k}\|^2 &\leq \|A\|^2 \|(X - P_0 Y)\|^2 \\ &= \sigma_1^2(A) \mathbf{d}([X], [Y])^2. \end{aligned}$$

Therefore, we conclude that $\sigma_1(A)$ is an upper Lipschitz constant for map β_A . We still need to show that this bound is achieved (i.e., it is optimal). For X_{\max} and Y_{\max} defined in part 1) of theorem 10,

$$\|\beta_A(X_{\max}) - \beta_A(Y_{\max})\|^2 = \|\beta_A(X_{\max})\|^2 = \sum_{k=1}^D \langle u, a_k \rangle^2 = \sigma_1^2(A).$$

and $\mathbf{d}(X_{\max}, Y_{\max}) = 1$. Thus B_0 is the

optimal Lipschitz constant both for α_A and for β_A .

2) Lower Lipschitz constants.

(i) Let $x, y \in \mathbb{R}^d$ and define the auxiliary set

$$S = S(x, y) := \{j \in [D] : |\langle x - y, a_j \rangle| \leq |\langle x + y, a_j \rangle|\}$$

Then

$$\begin{aligned} \|\alpha(x) - \alpha(y)\|^2 &= \sum_{i=1}^D \left| |\langle a_i, x \rangle| - |\langle a_i, y \rangle| \right|^2 = \\ &= \sum_{i \in S} |\langle a_i, x - y \rangle|^2 + \sum_{i \in S^c} |\langle a_i, x + y \rangle|^2 \geq \\ &\sigma_d^2(A[S]) + \sigma_d^2(A[S^c]) \mathbf{d}_{\text{PR}}([x], [y])^2 \geq A_0^2 \mathbf{d}_{\text{PR}}([x], [y])^2. \end{aligned}$$

So A_0 is a lower Lipschitz bound for α_A , but we still need to show that it is optimal.

Let I_0 be the optimal partition, and let u_1, u_2 be normalized left singular vectors as in the statement of Theorem 10. Then:

$$\begin{aligned} \|\alpha_A(u_1 + u_2) - \alpha_A(u_1 - u_2)\|^2 &= \\ &= \sum_{i=1}^D \left| |\langle a_i, u_1 + u_2 \rangle| - |\langle a_i, u_1 - u_2 \rangle| \right|^2 = \\ &= \sum_{i=1}^D \min(|\langle a_i, 2u_2 \rangle|^2, |\langle a_i, 2u_1 \rangle|^2) \leq \\ &\leq 4 \left(\sum_{i \in I_0} |\langle a_i, u_1 \rangle|^2 + \sum_{i \in I_0^c} |\langle a_i, u_2 \rangle|^2 \right) \\ &= 4(\sigma_d^2(A[I_0]) + \sigma_d^2(A[I_0^c])) = A_0^2 \mathbf{d}_{\text{PR}}([u_1 + u_2], [u_1 - u_2])^2, \end{aligned}$$

where we used again that $||a| - |b|| = \min(|a - b|, |a + b|)$ for any two real numbers $a, b \in \mathbb{R}$, and, for the inequality, at every $i \in [D]$ we made a choice between the two terms. Since the reverse inequality is also true, it follows that $x_{\min} = u_1 + u_2$ and $y_{\min} = u_1 - u_2$ achieve the lower bound A_0 for α_A .

(ii) Consider now the map β_A . Let $X, Y \in \mathbb{R}^{2 \times d}$ and define the auxiliary set

$$S = S(X, Y) := \{j \in [D] : |\langle x_1 - x_2 - y_1 + y_2, a_j \rangle| \leq |\langle x_1 - x_2 + y_1 - y_2, a_j \rangle|\}$$

Then, using Equation (5) we have that

$$\begin{aligned} \|\beta_A(X) - \beta_A(Y)\|^2 &= \\ \frac{1}{2} (\|\alpha_A(x_1 - x_2) - \alpha_A(y_1 - y_2)\|^2 + \|T_A(x_1 + x_2 - y_1 - y_2)\|^2) &= \\ \frac{1}{2} \sum_{j \in S} |\langle x_1 - x_2 - y_1 + y_2, a_j \rangle|^2 + |\langle x_1 + x_2 - y_1 - y_2, a_j \rangle|^2 + \\ + \frac{1}{2} \sum_{j \in S^c} |\langle x_1 - x_2 + y_1 - y_2, a_j \rangle|^2 + |\langle x_1 + x_2 - y_1 - y_2, a_j \rangle|^2 &= \\ = \sum_{j \in S} |\langle x_1 - y_1, a_j \rangle|^2 + |\langle x_2 - y_2, a_j \rangle|^2 + \\ + \sum_{j \in S^c} |\langle x_1 - y_2, a_j \rangle|^2 + |\langle x_2 - y_1, a_j \rangle|^2 \geq \\ \geq \sigma_d^2(A[S]) (\|x_1 - y_1\|^2 + \|x_2 - y_2\|^2) + \end{aligned}$$

$$\begin{aligned} + \sigma_d^2(A[S^c]) (\|x_1 - y_2\|^2 + \|x_2 - y_1\|^2) &\geq \\ \geq A_0^2 \mathbf{d}([X], [Y])^2. \end{aligned}$$

Therefore A_0 is a lower Lipschitz constant for β_A .

It remained to prove that this bound is tight, i.e., it is achieved. Let X_{\min} and Y_{\min} be as in the statement of Theorem 10. Then

$$\begin{aligned} \|\beta_A(X_{\min}) - \beta_A(Y_{\min})\|^2 &= \\ \frac{1}{2} (\|\alpha_A(u_1 + u_2) - \alpha_A(u_1 - u_2)\|^2 + \|T_A(u_1 + u_2 - u_1 - u_2)\|^2) &= \\ \frac{1}{2} (\|\alpha_A(u_1 + u_2) - \alpha_A(u_1 - u_2)\|^2) = A_0^2 \mathbf{d}([X_{\min}], [Y_{\min}])^2 \end{aligned}$$

where the last equality follows from the fact that the lower Lipschitz constant of α_A is achieved by $u_1 + u_2$ and $u_1 - u_2$, and the fact that $\mathbf{d}([X_{\min}], [Y_{\min}])^2 = 2$.

So A_0 is indeed the optimal lower Lipschitz constant for β_A . \square

III. CONCLUSION

In this paper we analyzed two representation problems, one arising in the phase retrieval problem and the other one in the context of permutation invariant representations. We showed that the real phase retrieval problem in a finite dimensional vector space H is entirely equivalent to the permutation invariant representations for the space $V = \mathbb{R}^{2 \times \dim(H)}$. Our analysis proved that phase retrievability is equivalent to the universal key property in the case of encoding $2 \times d$ matrices. This result is derived based on the lattice space structure $(\mathbb{R}, +, \min, \max)$. It is still an open problem to understand the relationship between α_A and β_A in the case $n > 2$. A related problem is the implementation of the sorting operator using a neural network that has ReLU as activation function (or, even the absolute value $|\cdot|$). Efficient implementations of such operator may yield novel relationships between α_A and β_A , in the case $n \geq 3$.

ACKNOWLEDGMENT

The authors have been supported in part by a NSF award under grant DMS-2108900 and by the Simons Foundation.

REFERENCES

- [1] N. Kasai, M. Kakudo, and Kasai-Kakudo..., *X-ray Diffraction by Macromolecules*. Springer, 2005, vol. 80.
- [2] R. W. Gerchberg, "A practical algorithm for the determination of plane from image and diffraction pictures," *Optik*, vol. 35, no. 2, pp. 237–246, 1972.
- [3] J. R. Fienup, "Phase retrieval algorithms: a comparison," *Applied optics*, vol. 21, no. 15, pp. 2758–2769, 1982.
- [4] G. Zheng, *Fourier ptychographic imaging: A MATLAB® tutorial*. Morgan & Claypool Publishers, 2016.
- [5] N. Sissouno, F. Boßmann, F. Filbir, M. Iwen, M. Kahnt, R. Saab, C. Schroer, and W. zu Castell, "A direct solver for the phase retrieval problem in ptychographic imaging," *Mathematics and Computers in Simulation*, vol. 176, pp. 292–300, 2020.
- [6] A. Oppenheim, J. Lim, G. Kopec, and S. Pohlig, "Phase in speech and pictures," in *ICASSP'79. IEEE International Conference on Acoustics, Speech, and Signal Processing*, vol. 4. IEEE, 1979, pp. 632–637.
- [7] R. Balan, "On signal reconstruction from its spectrogram," in *2010 44th Annual Conference on Information Sciences and Systems (CISS)*. IEEE, 2010, pp. 1–4.
- [8] P. J. Winzer, "Modulation and multiplexing in optical communications," in *Conference on Lasers and Electro-Optics*. Optica Publishing Group, 2009, p. CTuL3.
- [9] P. Delsarte, J.-M. Goethals, and J. J. Seidel, "Spherical codes and designs," in *Geometry and Combinatorics*. Elsevier, 1991, pp. 68–93.
- [10] R. Balan, B. G. Bodmann, P. G. Casazza, and D. Edidin, "Painless reconstruction from magnitudes of frame coefficients," *Journal of Fourier Analysis and Applications*, vol. 15, no. 4, pp. 488–501, 2009.
- [11] T. Heinosaari, L. Mazzarella, and M. M. Wolf, "Quantum tomography under prior information," *Communications in Mathematical Physics*, vol. 318, no. 2, pp. 355–374, 2013.
- [12] M. Kech and M. M. Wolf, "Quantum tomography of semi-algebraic sets with constrained measurements. preprint," *ArXiv*, vol. 1507, 2015.
- [13] E. J. Candes, T. Strohmer, and V. Voroninski, "Phaselift: Exact and stable signal recovery from magnitude measurements via convex programming," *Communications on Pure and Applied Mathematics*, vol. 66, no. 8, pp. 1241–1274, 2013.
- [14] E. J. Candes, Y. C. Eldar, T. Strohmer, and V. Voroninski, "Phase retrieval via matrix completion," *SIAM review*, vol. 57, no. 2, pp. 225–251, 2015.
- [15] X. Li and V. Voroninski, "Sparse signal recovery from quadratic measurements via convex programming," *SIAM Journal on Mathematical Analysis*, vol. 45, no. 5, pp. 3019–3033, 2013.
- [16] E. J. Candès and X. Li, "Solving quadratic equations via phaselift when there are about as many equations as unknowns," *Foundations of Computational Mathematics*, vol. 14, pp. 1017–1026, 2014.
- [17] A. Conca, D. Edidin, M. Hering, and C. Vinzant, "An algebraic characterization of injectivity in phase retrieval," *Applied and Computational Harmonic Analysis*, vol. 38, no. 2, pp. 346–356, 2015.
- [18] S. Shamma, "Analysis of speech dynamics in the auditory system," *Dynamics of Speech Production and Perception: Life and Behavioural Sciences*, pp. 335–342, 2006.
- [19] J. Li, Y. Xie, P. Stoica, X. Zheng, and J. Ward, "Beampattern synthesis via a matrix approach for signal power estimation," *IEEE Transactions on Signal Processing*, vol. 55, no. 12, pp. 5643–5657, 2007.
- [20] G. S. Thakur, "Reconstruction of bandlimited functions from unsigned samples," *Journal of Fourier Analysis and Applications*, vol. 17, pp. 720–732, 2010.
- [21] Y. Chen, C. Cheng, Q. Sun, and H. Wang, "Phase retrieval of real-valued signals in a shift-invariant space," *Applied and Computational Harmonic Analysis*, vol. 49, no. 1, pp. 56–73, 2020.
- [22] P. Jaming, "Phase retrieval techniques for radar ambiguity problems," *Journal of Fourier Analysis and Applications*, vol. 5, pp. 309–329, 1999.
- [23] —, "Uniqueness results for the phase retrieval problem of fractional fourier transforms of variable order," *arXiv preprint arXiv:1009.3418*, 2010.
- [24] S. Mallat, "Group invariant scattering," *Communications on Pure and Applied Mathematics*, vol. 65, no. 10, pp. 1331–1398, 2012.
- [25] J. Bruna and S. Mallat, "Invariant scattering convolution networks," *IEEE transactions on pattern analysis and machine intelligence*, vol. 35, no. 8, pp. 1872–1886, 2013.
- [26] S. Bahmani and J. Romberg, "Phase retrieval meets statistical learning theory: A flexible convex relaxation," in *Artificial Intelligence and Statistics*. PMLR, 2017, pp. 252–260.
- [27] R. Balan, P. Casazza, and D. Edidin, "On signal reconstruction without noisy phase," *Appl. Comput. Harmon. Anal.*, vol. 20, pp. 345–356, 2006.
- [28] A.S. Bandeira, J. Cahill, D. Mixon, A.A. Nelson, "Saving phase: Injectivity and stability for phase retrieval," *Appl. Comp. Harm. Anal.*, vol. 37, no. 1, pp. 106–125, 2014.
- [29] A. H. J. Cahill, A. Contreras, "Complete set of translation invariant measurements with lipschitz bounds," *Appl. Comput. Harm. Anal.*, vol. 49, no. 2, pp. 521–539, 2020.
- [30] R. Balan, N. Haghani, and M. Singh, "Permutation invariant representations with applications to graph deep learning," 2022.
- [31] J. Cahill, J. W. Iverson, D. G. Mixon, and D. Packer, "Group-invariant max filtering," *arXiv:2205.14039 [cs.IT]*, pp. 1–35, 2022.
- [32] N. Dym and S. J. Gortler, "Low dimensional invariant embeddings for universal geometric learning," *arXiv:2205.02956 [cs.LG]*, pp. 1–23, 2022.
- [33] R. Balan and Y. Wang, "Invertibility and robustness of phaseless reconstruction," *Applied and Comput. Harmon. Analysis*, vol. 38, no. 3, pp. 469–488, 2015.
- [34] R. Balan and D. Zou, "On lipschitz analysis and lipschitz synthesis for the phase retrieval problem," *Linear Algebra and Applications*, vol. 496, pp. 152–181, 2016.